

Microbiology Research Journal International

Volume 34, Issue 12, Page 149-161, 2024; Article no.MRJI.127292 ISSN: 2456-7043, NLM ID: 101726596 (Past name: British Microbiology Research Journal, Past ISSN: 2231-0886, NLM ID: 101608140)

Nutritional, Antioxidant and Antimicrobial Properties of Citrus Peels: A Sustainable Valorization Approach

Sudeshna Sengupta ^a, Rojina Khatun ^a, Aanchal Kumari ^a, Roshlina Rai ^a, Subarna Dutta ^a, Torsha Mondal ^a, Banani Chattaraj ^a and Malavika Bhattacharya ^{a*}

^a Department of Biotechnology, Techno India University, EM-4, Sector-V, Salt Lake, Kolkata, West Bengal-700091, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: https://doi.org/10.9734/mrji/2024/v34i121517

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/127292

Original Research Article

Received: 08/10/2024 Accepted: 10/12/2024 Published: 18/12/2024

ABSTRACT

Background: Citrus peels are often discarded as waste despite being rich in bioactive compounds with potential health benefits. Traditional extraction methods for these compounds are not always sustainable and can lead to environmental harm. Using a sustainable vaporization approach, the bioactive compounds in citrus peels can be efficiently extracted and retained for various applications. Valorization strategies have been used for various waste streams, including agro-industrial by-products and olive mill wastewater.

Cite as: Sengupta, Sudeshna, Rojina Khatun, Aanchal Kumari, Roshlina Rai, Subarna Dutta, Torsha Mondal, Banani Chattaraj, and Malavika Bhattacharya. 2024. "Nutritional, Antioxidant and Antimicrobial Properties of Citrus Peels: A Sustainable Valorization Approach". Microbiology Research Journal International 34 (12):149-61. https://doi.org/10.9734/mrji/2024/v34i121517.

^{*}Corresponding author: E-mail: malavikab@gmail.com;

Aims: Using a sustainable vaporization method, this study seeks to examine the antibacterial, antioxidant, and nutritional qualities of citrus peels.

Methods: waste valorization remains a promising approach for sustainable waste management and resource recovery. Quantitative studies and antimicrobial activities were carried out using peels of C. aurantifolia (Key Lime), C. maxima (Pomelo), C. sinensis (orange), Citrus limetta (Sweet Lemon), and C. reticulata. Citrus aurantiifolia is believed to be a hybrid of Citrus medica, Citrus grandis, and a Micro-citrus species, with significant antibacterial activity.

Result: *Citrus limetta* Mosambi is gaining popularity due to its delicious taste and high vitamin-C content. *Citrus sinensis* is a small, spherical fruit rich in vitamin C and vitamin A. Pummelo, or *Citrus maxima*, is known for its large, sweet fruits with abortifacient and menstrual stimulant properties. Peels of *C. maxima* exhibit the highest amount of antioxidant activity equivalent to ascorbic acid and display significant antimicrobial activity against *Klebsiella species*. Citrus reticulata, found mainly in eastern India, has a maximum amount vitamin C among citrus species and exhibits high antimicrobial activity.

Keywords: Valorization; citrus fruits; vitamin-C; antibacterial; agro-industrial waste; nutraceutical.

1. INTRODUCTION

Citrus fruits play a common role in our daily lives, and their peels, pulps, and seeds offer various benefits. Since childhood, we've known that citrus fruits are rich in vitamins and are good for human health. The peels of fruits make up a significant portion of their weight and are often discarded as waste. While the pharmaceutical industry and synthetic medications are widely relied upon in the modern world, many people are unaware of the beneficial qualities of numerous herbs and vegetables. One of the main antioxidants in circulation, vitamin C, has immune-stimulating anti-inflammatory and properties and is necessary for optimal health at every stage of life. Scientific pathology may recommend synthetic pharmaceuticals to treat a variety of ailments, but it's important to focus on our natural healing process, which is something that Mother Nature always does her best to support (Sengupta et al. 2023). However, it's worth noting that these fruit remnants can still be utilized in several ways rather than simply being thrown away. Citrus fruits are known for being a rich source of phytochemicals and biologically active compounds that contribute to their healthpromoting properties. It's recommended to incorporate a variety of fruits and vegetables into your diet to ensure a well-rounded intake of nutrients and bioactive compounds. Valorization is a concept aimed at sustainably managing industrial residues by recovering valuable components or producing useful products from waste materials (Federici et al. 2009). It encompasses various approaches, including physical, chemical, and biological methods, to transform waste into economically valuable and environmentally friendly products (Faheem et

al.2021). The concept aligns with industrial ecology principles, promoting loop-closing and resource efficiency (Nzihou & Lifset 2010). Valorization strategies have been applied to diverse waste streams, such as agro-industrial by-products and olive mill wastewaters, focusing on extracting antioxidants and producing enzymes. However, the valuation of knowledge in the valorization process presents challenges due to its subjective nature. Financial valuation methods, particularly the income approach, are commonly used but require assumptions that may be difficult to validate (Andriessen 2005). Despite these challenges, waste valorization remains a promising approach for sustainable waste management and resource recovery.

When citrus peel is used as a substrate in anaerobic digestion processes, it can inhibit the process. This is because the high content of certain compounds in citrus peel, such as dlimonene, can interfere with the activity of methanogenic bacteria responsible for methane production. As a result, this interference can lead to reduced biogas production and longer lag phases before methane production begins (Fagbohungbe et al. 2016).

Citrus fruit peels contain high levels of flavonoids, polyphenols, and dietary fiber. These components have antioxidant, anti-inflammatory, and potential anti-cancer properties. Extracts from citrus peels have been studied for their cholesterol-lowering effects, skin health benefits, and as potential natural remedies for various health conditions. Fruit residues, often considered waste, have the potential to be valuable nutraceutical resources. Citrus peels, pulp, and seeds contain various beneficial compounds that offer health-promoting properties. By effectively utilizing these residues, we can reduce waste and harness their nutraceutical potential (Rafig et al. 2018). The value of citrus fruit residues as nutraceuticals can be recognized by finding innovative ways to use them, thus reducing waste and benefiting from their health-promoting properties. Proper processing and guality control are essential to maximize their safety and effectiveness as nutraceutical resources. The use of citrus peel waste in the medicine industry involves extraction techniques to isolate and concentrate the bioactive compounds. Ongoing research and development aim to explore their therapeutic potential, optimize extraction methods, and develop innovative drug delivery systems. While citrus peel and its bioactive compounds show promise in medicine, it is crucial to ensure appropriate quality control, standardization, and safety evaluation when using them for medicinal purposes (Baker et al. 2022).

Citrus peels were once considered waste, but they are now valued for their medicinal properties and are used in various applications, including food products. Citrus peels are rich in vitamins, fibers. and bioactive compounds, which contribute to their potential health benefits. Their antioxidant properties help to eliminate harmful free radicals in the body, providing protection against degenerative conditions. They also have antimicrobial properties that can inhibit the growth of certain bacteria and fungi. Additionally, researchers have taken note of their antiinflammatory and anticancer properties. Whether in the form of powder or essential oils, citrus peels have been successfully incorporated into food products to enhance their nutritional value without compromising taste and sensorv attributes when used in appropriate amounts. The journey of citrus peels, from being discarded as waste to being utilized in the laboratory and eventually making their way to the dining table, is a testament to their versatility and potential for both health and economic benefits (Ademosun 2022).

The utilization of waste materials generated from citrus fruit processing, such as peels and pulp, for the extraction of cellulose and nanocellulose, is a fascinating area of research. Cellulose is a natural polymer found in plant cell walls and has various applications due to its unique properties. In the present study, the waste materials from citrus were treated with an alkaline solution to extract cellulose. The cellulose was then

subjected to acid hydrolysis, resulting in the reduction of fiber size and the formation of nanocellulose. X-ray diffraction analysis revealed that the cellulose had an amorphous structure, indicating the breakdown of the crystalline regions during the extraction process. At high magnification, break points in the cellulose fibers were observed, resembling carbon nanotubes in appearance. This indicates that the acid treatment caused structural modifications to the cellulose fibers. Overall, this study highlights the potential of citrus peel waste as a valuable source of cellulose and nanocellulose. By extracting and modifying cellulose from citrus waste, researchers have discovered a novel application for this abundant agricultural byproduct. The utilization of citrus peel waste for cellulose extraction offers both environmental and economic benefits, as it reduces waste generation and provides a sustainable alternative to traditional cellulose sources (Naz et al. 2016). Citrus fruits generate significant waste during processing, including peels, seeds, and pomace. Proper management and utilization of this waste minimize environmental impacts and can resource utilization for optimize a more sustainable approach to citrus fruit processing.

2. MATERIALS AND METHODS

2.1 Samples Collection and Preparation

Peels of the citrus species thatwere taken for the experiments were *Citrus aurantifolia*, *Citrus limetta*, *Citrus reticulata* (Aroma King lemon/gandhoraj), *Citrus sinensis*, and *Citrus maxima*.

Peels of every citrus species have been attached for a clear view [Fig. 1].

2.2 Quantitative Nutritional Analysis

Total vitamin C and protein estimation by UV spectrophotometric method, followed by vitamin C quantification assay using the potassium iodate and potassium iodine titration method (Pathy 2018). Bradford assay, and Lowry assay (Bradford 1976). Carbohydrate estimation was done through the Anthrone method (Yemm & Willis 1954).

2.3 In vitro Antioxidant Assay Using DPPH

The stability of the extracts' DPPH (2, 2-diphenyl-2-picrylhydrazyl) free radical scavenging activity was assessed using the usual method (Shen et al. 2010).

Sengupta et al.; Microbiol. Res. J. Int., vol. 34, no. 12, pp. 149-161, 2024; Article no.MRJI.127292



Citrus sinensis (Orange)

Citrus limetta (Sweet lemon)



Citrus maxima (pomelo)

Citrus reticulata (Gandhoraj lemon)



Citrus aurantifolia (Keylime)

Fig. 1. These samples were taken

2.4 In vitro Antimicrobial Assay

The antibacterial assay was performed using the disc diffusion method against Gram-negative *Klebsiella sp. & Enterobacter sp.* and Gram-positive bacteria *Staphylococcus epidermis*, *Bacillus sp.* The bacteria strains were maintained in 100 mL of nutrient broth at 37°C and incubated overnight.

3. RESULTS

3.1 Quantitative Nutritional Analysis of Peels of the Samples

Out of five samples, *C. reticulata* (10.85 \pm 2.49 mg/ml), *C. lime* (10.46 \pm 2 mg/ml), and *C. sinensis* (8.96 \pm 1.6 mg/ml) have a higher amount of vitamin *C* than the other samples. Sd was measured (n = 3) for each of the samples.



Fig. 2. Vitamin C amount in the peel samples



Fig. 3. Protein amount in the peel of the samples

Peels of the citrus fruits have enough protein, out of these *C. sinensis* has a maximum amount of protein 422.76±8.76 (mg BSAE eq. /g FW) median *C. reticulata* 231.7±4.0115 (n=3) for each of the samples [Fig. 3].

The result clearly shows *C. sinensis* maximum carbohydrate of 2643.08 ± 4.98 equivalent to glucose, and a median of 905.95 ± 6.47 *C. limetta* Sd measured (n=3) for each of the samples [Fig. 4].

C. maxima and C. sinensis have maximum antioxidant activity equivalent to ascorbic acid than others. Sd measured (n=3) for each of the samples [Fig. 5].

"SD measured (n=3)" is mentioned throughout the manuscript and suggests that the experiment was conducted three times, it indicates that the standard deviation was calculated based on data obtained from three experimental runs.



Sengupta et al.; Microbiol. Res. J. Int., vol. 34, no. 12, pp. 149-161, 2024; Article no.MRJI.127292

Fig. 4. Carbohydrate amount in the peel samples



Fig. 5. Antioxidant activity of the peel samples

The result clearly shows that all of the samples have a similar amount of antimicrobial activity among other samples. ampicillin was the control, and Sd was measured (n=3) for each sample [Fig 6a].

The results indicate that C. limetta and C. reticulata exhibit similar levels of antimicrobial activity compared to other samples. Ampicillin was used as the control, and each sample's

standard deviation (SD) was measured (n=3) [Fig. 7a-7b].

The result clearly shows among all the samples C. reticulata, C. aurantiifolia, and C. sinensis showed maximum antimicrobial activity among other samples., ampicillin was the control, Sd measured (n=3) for each of the samples [Fig. 8a-8b].



Sengupta et al.; Microbiol. Res. J. Int., vol. 34, no. 12, pp. 149-161, 2024; Article no.MRJI.127292

Fig. 6a. Diagrammatic representation of the inhibitory zones measured in mm



Fig. 6b. The measurement area is the disc diffusion results using the gram (-) bacteria Enterobacter sp. inhibitory zones (mm scale)



Sengupta et al.; Microbiol. Res. J. Int., vol. 34, no. 12, pp. 149-161, 2024; Article no.MRJI.127292

Fig. 7a. Diagrammatic representation of the inhibitory zones measured in mm Ampicillin control



Fig. 7b. Shows the disc diffusion results using the gram (+) bacteria *Bacillus sp.* inhibitory zones (mm scale) are the area of measurement



Sengupta et al.; Microbiol. Res. J. Int., vol. 34, no. 12, pp. 149-161, 2024; Article no.MRJI.127292

Fig. 8a. The diagrammatic representation of the inhibitory zones measured in mm



Fig. 8b. Shows the disc diffusion results using the gram (+) bacteria *Staphylococcus* epidermis inhibitory zones (mm scale) are the area of measurement



Sengupta et al.; Microbiol. Res. J. Int., vol. 34, no. 12, pp. 149-161, 2024; Article no.MRJI.127292

Fig. 9a. The diagrammatic representation of the inhibitory zones measured in mm .



Fig. 9b. Shows the disc diffusion results using the gram (-) bacteria *Klebsiella* inhibitory zones (mm scale) are the area of measurement

The result clearly shows that *C. limetta and C. maxima* showed maximum antimicrobial activity among other samples. ampicillin was the control, and Sd measured (n=3) for each of the samples [Fig. 9a-9b].

4. DISCUSSION

In recent years, research has demonstrated connections between oral microbiota and systemic illnesses, including head and neck cancer (Amon et al. 2019). Ampicillin is a commonly used antibiotic that is effective against infections caused by various susceptible organisms, such as intestinal bacilli, salmonella, shigella, enterococci, listeria, and some forms of hemophilic bacilli. Citrus fruits are popular worldwide and offer various nutritional and medicinal benefits. For example. Citrus aurantiifolia is believed to be a hybrid of Citrus medica, Citrus grandis, and a Micro-citrus species. The spherical fruit, with a diameter of 25-50 mm, is typically harvested when green when ripened. and turns yellow Citrus aurantiifolia exhibits significant antibacterial activity. Citrus aurantiifolia and C. maxima show antimicrobial the hiahest activity. Citrus reticulata, also known as the Aroma King Lemon or Gandhoraj, combines the flavors of mandarin orange and lime. This species is found mainly in eastern India. The peel of Citrus reticulata has the highest amount of vitamin C among the five citrus species and exhibits high antimicrobial activity against Staphylococcus epidermis and Bacillus species. Citrus limetta Mosambi, a popular lime variety, is gaining worldwide popularity due to its delicious taste and high vitamin C content. Citrus sinensis, resembling a typical lime, is a small, spherical fruit rich in vitamin C and some vitamin A, making it highly nutritious. The peel and pulp of Citrus sinensis have the maximum amount of protein among all species, while the seeds and peel are good sources of carbohydrates.

Pummelo, or Citrus maxima, is a citrus tree producing large, sweet fruits with a history of use in herbal and traditional medicine due to its abortifacient and menstrual stimulant properties. Excessive use of these drugs can lead to severe liver damage, vomiting, and bleeding, which can be fatal. Contact with rue can also cause severe phytophotodermatitis and burn-like blisters after sun exposure. Peels of C. maxima exhibit the highest amount of antioxidant activity equivalent to ascorbic acid and display significant antimicrobial activity against Klebsiella species.

Citrus trash can provide nanocellulose, a flexible and biodegradable substance that provides an eco-friendly approach to waste management in the beverage sector (Fan et al. 2016, Mariño et al. 2015). The shape and characteristics of the final nanofibers are influenced by different extraction methods (Mariño et al. 2015). In comparison to conventional Hestrin-Schramm media, citrus peel and pomace enzymolysis (CPPE) medium has demonstrated a greater bacterial cellulose production (Fan et al. 2016). The isolated nanocellulose has average diameters of 10-60 nm and crystallinity indices between 55% and 65% (Fan et al. 2016, Mariño et al. 2015). Cellulose fiber size can be decreased by alkaline treatment followed by acid producina denselv hvdrolvsis. packed nanocellulose structures that resemble carbon nanotubes (Naz et al. 2016). In paleobotanical investigations, nitrocellulose, another cellulose derivative. has been employed: the characteristics of the resulting film are influenced by different solvents and plasticizers (Graham 1933).

Extracts from citrus peels have shown strong antibacterial action against a range of infections. Lemon peel extract shown encouraging results against Candida albicans, Escherichia coli, and Staphylococcus aureus (Ali et al. 2017). Likewise. Turkish citrus peel oils demonstrated potent antibacterial properties, with bergamot and lemon oils showing the most efficacy (Gülay et al. 2009). According to a comparative analysis, lemon essential oil had the most inhibitory effects against Candida albicans and both Gram-positive and Gram-negative bacteria (Alsakhawy et al. 2023). In citrus peel essential oils, limonene was found to be the main volatile component, although catechin and naringenin were important phenolic components (Alsakhawy et al. 2023). Citrus aurantium peel extract had exceptional qualities antifungal against Colletotrichum capsici, while Citrus sinensis peel extract indicated strong antibacterial activity, especially against Klebsiella pneumoniae (Madhuri et al. 2014).

5. CONCLUSION

Valuation techniques can have a significant impact on resource efficiency and waste reduction. For example, citrus peels can be repurposed to create valuable products such as animal feed, dietary supplements, and essential oils. The article effectively demonstrates how applying valuation methods to citrus fruit waste can highlight the financial and environmental benefits of implementing these strategies. Furthermore, it discusses the potential for scaling up these valuation methods and their support for environmentally friendly waste management techniques. This deeper investigation could enhance the overall impact of valuation and consider the potential for producing nutraceutical from citrus peels. These results demonstrate citrus waste's potential as a useful resource for the synthesis and use of nanocellulose also suggests that citrus peel extracts have potential as natural antimicrobial agents.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology.

Details of the AI usage are given below:

- 1. Elicit
- 2. Scispace
- 3. Grammarly

ACKNOWLEDGEMENTS

I would like to sincerely thank Techno India University for providing me with the knowledge and guidance required.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Ademosun, A. O. (2022). Citrus peels odyssey: From the waste bin to the lab bench to the dining table. *Applied Food Research*, 2(1), 100083. DOI: https://doi.org/10.1016/j.afres.2022.10008.
- Ali, J., Das, B., & Saikia, T. (2017). ANTIMICROBIAL ACTIVITY OF LEMON PEEL (CITRUS LIMON) EXTRACT. International Journal of Current Pharmaceutical Research, 9, 79-82.
- Alsakhawy, S.A., Baghdadi, H.H., El-shenawy, M., & El-Hosseiny, L.S. (2023). Comparative Phytochemical Composition

and Antimicrobial Activity of Citrus Peel Essential Oils and Phenolic Compounds. *Anti-Infective Agents*.

- Amon, S., Meier-Abt, F., Gillet, L. C., Dimitrieva, S., Theocharides, A. P., Manz, M. G., & Sensitive Aebersold, R. (2019). Quantitative Proteomics Human of Hematopoietic Stem and Progenitor Cells by Data-independent Acquisition Mass Spectrometry. Molecular R Cellular 18(7), 1454–1467. Proteomics. DOI: https://doi.org/10.1074/mcp.tir119.001431.
- Andriessen, D. (2005). Value, Valuation, and Valorisation.
- Baker, D. H. A., Ibrahim, E. A., & Salama, Z. a.
 E. R. (2022). Citrus Peels as a Source of Bioactive Compounds with Industrial and Therapeutic Applications. In *Biochemistry*. DOI:

https://doi.org/10.5772/intechopen.99591.

- Bradford M.M. (1976) Rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Anal. Biochem; 72:248–254.
- Fagbohungbe MO, Herbert BM, Hurst L, Li H, Usmani SQ, Semple KT. Impact of biochar on the anaerobic digestion of citrus peel waste. Bioresour Technol. 2016 Sep;216:142-9. doi: 10.1016/j.biortech.2016.04.106. Epub 2016 May 10. PMID: 27236401.
- Faheem, M., Hassan, M.A., Mehmood, T., Hashim, S., & Ashraf, M.A. (2021). Waste Valorization: Physical, Chemical, and Biological Routes. *Sustainable Solutions for Environmental Pollution*.
- Fan, X., Gao, Y., He, W., Hu, H., Tian, M., Wang, K., & Pan, S. (2016). Production of nano bacterial cellulose from beverage industrial waste of citrus peel and pomace using Komagataeibacter xylinus. *Carbohydrate polymers*, *151*, 1068-1072.
- Federici, F., Fava, F., Kalogerakis, N., &Mantzavinos, D. (2009). Valorisation of agro-industrial by-products, effluents and waste: concept, opportunities and the case of olive mill wastewaters. *Journal of Chemical Technology & Biotechnology, 84*, 895-900.
- Graham, R.J. (1933). Preparation of Palaeobotanical Sections by the Peel Method. *Biotechnic & Histochemistry, 8*, 65-68.
- Gülay, F.G., Tavman, A., & Türker, G. (2009). Antimicrobial activity of Turkish citrus peel oils.

- Madhuri, S., Hegde, A., Srilakshmi, N., & Kekuda, T.R. (2014). Antimicrobial activity of *Citrus sinensis* and *Citrus aurantium* peel extracts. *Journal of Pharmaceutical and Scientific Innovation*, 3, 366-368.
- Mariño, M.A., Lopes da Silva, L., Durán, N., & Tasić, L. (2015). Enhanced Materials from Nature: Nanocellulose from Citrus Waste. *Molecules, 20*, 5908 - 5923.
- Naz, S., Ahmad, N., Akhtar, J., Ahmad, N. M., Ali, A., & Zia, M. (2016). Management of citrus waste by switching in the production of nanocellulose. *IET Nanobiotechnology*, *10*(6), 395–399. DOI: https://doi.org/10.1049/iet-nbt.2015.0116.
- Naz, S., Ahmad, N., Akhtar, J., Ahmad, N.M., Ali, A., & Zia, M. (2016). Management of citrus waste by switching in the production of nanocellulose. *IET nanobiotechnology*, 10 6, 395-399.
- Nzihou, A., &Lifset, R. (2010). Waste Valorization, Loop-Closing, and Industrial Ecology. *Journal of Industrial Ecology, 14*.
- Pathy, Krishna Sarma. (2018). Process for Preparation of Vitamin C and Method for Determination of Vitamin C in Tablets Volume 1 - Issue 3 Krishnasarmapathy*.

- Rafiq, S., Kaul, R., Sofi, S., Bashir, N., Nazir, F., &Nayik, G. A. (2018). Citrus peel as a source of functional ingredient: A review. *Journal of the Saudi Society of Agricultural Sciences*, 17(4), 351–358. DOI: https://doi.org/10.1016/j.jssas.2016.07.006.
- Sengupta, S., Banerjee, S., Nayek, S. N., Das, P., Chakraborty, D., Mukherjee, A., Mandal, T., Majumder, A., Saha, M., Chatterjee, S., & Bhattacharya, M. (2023).
 A brief comparative study of the natural sources (lemons) in the basis of protein, vitamin C, their antibacterial, anthelminthic and cell viability on immune cells. *International Journal of Herbal Medicine*, *11*(5), 14–21. DOI: https://doi.org/10.22271/flora.2023.v11.i5a. 8831.
- Shen Q, Zhang B, Xu R, Wang Y, Ding X, Li P. (2010). Antioxidant activity In vitro of Selenium-contained protein from the Seenriched Bifidobacteriu manimalis 01. Anaerobe. 16.
- Yemm, E. W., & Willis, A. J. (1954). The estimation of carbohydrates in plant extracts by anthrone. *Biochemical Journal*, *57*(3), 508–514. DOI: https://doi.org/10.1042/bj0570508.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/127292