

Journal of Experimental Agriculture International

Volume 46, Issue 12, Page 581-588, 2024; Article no.JEAI.127684 ISSN: 2457-0591 (Past name:American Journal of Experimental Agriculture,Past ISSN: 2231-0606)

# Effect of Genotypes, Fertilizer Application & Crop Geometry on Growth, Yield and Yield Attributes in Coriander (*Coriandrum sativum* L.)

# R. C. Aswani<sup>a\*</sup>, G. S. Gathiye<sup>b</sup> and S. K. Tyagi<sup>b</sup>

<sup>a</sup> Directorate of Extension Services, RVSKVV, Gwalior (M. P.), India. <sup>b</sup> KVK, Dhar and Khargone, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: https://doi.org/10.9734/jeai/2024/v46i123164

#### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/127684

**Original Research Article** 

Received: 12/10/2024 Accepted: 18/12/2024 Published: 27/12/2024

# ABSTRACT

A field experiment was conducted during rabi 2019-20 & 2020-21 at KNK Horticulture College. Treatments comprised of three crop geometry viz., 20x10 cm (S1), 25x10 cm (S2) and 30x10cm (S3), two fertilizer levels viz., F145:35:35:20 N:P:K:S ha<sup>-1</sup>, F260:45:45:30 N:P:K:S ha<sup>-1</sup> and three genotypes viz., Rcr 41 (V1), Ajmer Coriander ACr<sup>-2</sup> (V2) and Pant Haritima as a check (V3). The experiment was laidout in Randomized Block Design (RBD) with factorial concept. Result revealed that application of N:P:K:S 60:45:45:30 (F2) Kg<sup>-1</sup> ha with a row spacing of 30x10 cm (S3) significantly improved plant height, number of branches/plant, while minimum days to 50% flowering was recorded in 20x10cm (S1), whereas the highest number of umbels/plant, test weight (g), Umbellates Plant-1, seed yield (kg ha-1), straw yield (kg ha-1), biological yield (kg ha-1),

\*Corresponding author: E-mail: rameshaswani69@gmail.com;

*Cite as:* Aswani, R. C., G. S. Gathiye, and S. K. Tyagi. 2024. "Effect of Genotypes, Fertilizer Application & Crop Geometry on Growth, Yield and Yield Attributes in Coriander (Coriandrum Sativum L.)". Journal of Experimental Agriculture International 46 (12):581-88. https://doi.org/10.9734/jeai/2024/v46i123164.

harvest index (%) was recorded in 30x10 cm(S3) Highest No. of nodes (11.20, 11.08 & 11.28) at harvest in S3, F2and V2and primary branches were recorded (8.86, 8.65 & 8.58) in S3, F2 and V2 respectively for growth parameters whereas, maximum No. of umbels and No. of umbellate (39.26,38.20,42.39&139.91,134.66,140.82) was recorded at harvest inS3,F2and V2 respectively. Maximum Seed yield (13.22, 12.62, 12.80 & 25.77, 24.57, 24.76) was recorded in S3, F2and V2 respectively (kg ha-1). Ajmer Coriander ACr-1 (V2) recorded an additional net returns of Rs. 89273 and Rs. 82510 over Pant Haritima (V3) and Rcr 41 (V1). Highest B:C ratio was obtained with Ajmer Coriander ACr-1(3.36) as compared to Pant Haritima (3.34) and Rcr 41 (2.93) respectively.

Keywords: Coriander; genotypes; umbel; B:C ratio; crop geometry.

### 1. INTRODUCTION

"Coriander (Coriandrum sativum L.) is one of the most important an annual spice, condiment, medicinal and wellknown herb for it's mild pleasant aroma and test belongs to the family Apiaceae. Coriander plant gives two primary products that are used for flavouring, beverages and seeds are being used for preparing value added products such as curry powder, dhana dal, condiment and essential oil oleoresin. The essential oil content of locally available seeds of coriander is 0.2-0.3 per cent whereas major chemical constituent of essential oil are dlinalool, linalyl acetate along with other important constituents such as thymol. geraniol, carophyllene and pinene. The green leaves are good source of vitamin C (12.0 mg) and (175 IU/100 g), while seed are rich in carbohydrates and protein content. India is the largest producer, consumer and exporter of coriander in the world. In India the major coriander growing states are Rajasthan, Gujarat, Andhra Pradesh, Tamil Nadu, Karnataka, Madhya Pradesh, Bihar, Uttar Pradesh (Moosavi et al., 2012). The area under coriander cultivation in India is 663 thousand hectare and production is 609 thousand MT and in Madhya Pradesh, coriander is cultivated over an area of 224 thousand hectare and production is 116.27 thousand MT(Horticulture (3rd Adv. Estimate) 2023-24, Deptt. of Ag.& FW, Gol). In the state of Madhya Pradesh coriander is mostly grown in Mandsaur, Neemuch, Shajapur, Rajgarh, Guna and Vidisha. Row spacing is an important factor for better growth and yield of the plant (Mishra et al., 2019). Optimum number of plants is required per unit area to utilize efficiently the available production factors such as water, nutrient, light and CO2. Maximum exploitation of these factors is achieved when the plant population puts forth maximum pressure on all the factors of production" Narukaet al. (2012). "Nitrogen is a constituent of proteins, enzymes, hormones, vitamins, alkaloids, chlorophyll etc. Adequate supply of N promotes higher

photosynthetic activity and vigorous vegetative growth and as a result, the plants turn in to dark green colour(Painkaraet al., 2024). A high N supply favours the conversion of carbohydrate into protein which, in turn promotes the formation of protoplasm. Protoplasm, being highly hydrated, is conducive for the succulent plant growth" (Mehta et al. 2013).

### 2. MATERIAL AND METHODS

A field experiment was conducted during rabi 2019-20 & 2020-21 at KNK Horticulture College, experimental Research Farm, PatanMandsaur. The experiment was laid out in factorial randomized block design having three replications. There were eiaht treatment combinations, consisted of three row spacing viz., 20x10 cm, 25x10 cm and 30x10 cm and two fertilizer levels viz., F1-45:35:35:20 N:P:K:S ha-1, F2- 60:45:45:30 N:P:K:S ha-1. Seeds were sown on 24th& 26thOctober, 2019-20 & 2020-21 in furrows at different row spacings. The seed of three genotypes viz., Rcr 41 (V1), Ajmer Coriander ACr<sup>-1</sup> (V2) and Pant Haritima as a check (V3) @15 kg/ha was used for sowing. The unit plot size was 2.0 m x3.0 m. The seeds (fruits) were rubbed for separating the mercers (seeds) and were also treated with Bavistin @ 2g kg-1 of seeds before sowing. First manual weeding was done at 30 days and second at 60 DAS to control the season bound weeds. after sowing (DAS). Plant thinning was also done with weeding to maintain the required plant population. The irrigation was done at 20 days in terval after emergence according to the need of the field soil. Data were collected from the inner rows of each plot to avoid the border effect. In each unit plot 5 plants were selected randomly for recording data on plant growth and yield parameters. Plot yield was converted to per hectare yield. The soil of the experimental field was sandy clay loam having low organic matter (0.27%), available nitrogen (224.5 kg ha-1), phosphorus (10.4 kg ha-1) and enough available

potassium (504.43kg ha<sup>-1</sup>), sulphur (9.06 kg ha<sup>-1</sup>) slightly saline alkali with pH (7.8) and EC (0.072dS m<sup>-1</sup>). Irrigation was applied immediately after sowing. All other standard package of practices were followed during whole cropping season. Seed yield, yield attributes and growth parameter were recorded in five plant of each plot and average was worked out. The collected data were properly analyzed statistically through method suggested by Panse and Sukhatme (1985).

#### Availability of macro & micro nutrients in an experimental soil

S.	Ν	Р	К	S	Zn	Cu	Fe	Mn
No.	(kg ha <sup>-1</sup> )	(mg kg⁻¹)	(mg kg <sup>-1</sup> )	(mg kg⁻¹)	(mg kg⁻¹)			
1.	224.5	10.4	504.43	9.06	0.36	3.68	4.72	5.16

#### 3. RESULTS AND DISCUSSION

Effect of crop geometry (row spacing): The results clearly revealed that row spacing as well as genotype significantly influenced different growth parameters (Table 1) viz., plant height, number of nodes and primary branches per plant and days taken to 50% flowering. Maximum plant height (cm) at 90 DAS (77.92) was attained at row spacing of  $20x10 \text{ cm} (S_1)$ . Significant increase in plant height from early stage of crop growth under closer row spacing seem to be due to mutual shading due of the dense population. This might have decreased the availability of light to the plants. The reduced light intensity at the base of the plant stem might have accelerated elongation of lower internodes resulting in plant height. Although, plant elongates rapidly due to mutual shading but beyond a certain level, elongation is checked due to reduced availability of photosynthates. These observations are in close conformity with finding of Nayak et al. (2009), Malav and Yadav (1997) and Kumar et. al (2006).

Treatments	P. height (cm)	P. height (cm)	No. of Nodes/	No. of Pri. branch	DTF (50%)
	at 90 DAS	at harvest	plant		
Crop geometry					
20x10 cm (S1)	77.92	82.45	8.94	6.98	60.62
25x10 cm (S <sub>2</sub> )	66.49	75.01	10.25	7.82	61.67
30x10cm (S <sub>3</sub> )	61.03	67.69	11.20	8.86	63.72
S. Em±	1.03	1.12	0.27	0.21	0.48
CD (P=0.05)	2.99	3.26	0.78	0.62	1.41
Fertilizer application					
N:P:K:S 45:35:35:20 (F <sub>1</sub> )	63.84	71.21	9.31	7.16	63.77
N:P:K:S 60:45:45:30 (F <sub>2</sub> )	73.14	78.94	11.08	8.65	60.29
S. Em±	0.84	0.91	0.22	0.17	0.39
CD (P=0.05)	2.44	2.66	0.63	0.50	1.15
Genotypes					
Rcr41 (V <sub>1</sub> )	63.08	68.74	9.67	7.45	61.45
Ajmer Coriander ACr <sup>-1</sup> (V <sub>2</sub> )	77.82	88.01	10.28	9.58	63.95
Pant Haritima (V <sub>3</sub> )	74.77	80.41	8.28	7.59	60.71
S. Em±	1.03	1.12	0.27	0.21	0.48
CD (P=0.05)	2.99	3.26	0.78	0.62	1.41
CV%	6.35	6.32	11.13	11.40	3.31

#### Table 1. Effect of spacing, fertilizer application and genotypes on different growth parameters of coriander

Treatments	No. of Umbels Plant <sup>-1</sup>	No. of Umbellates	Test Weight	Seed yield	Straw yield	<b>Biological yield</b>	Harvest
		Plant <sup>-1</sup>	(g)	(kg ha⁻¹)	(kg ha⁻¹)	(kg ha⁻¹)	index (%)
Crop geometry							
20x10 cm (S <sub>1</sub> )	35.38	121.31	6.64	10.59	22.80	33.38	31.78
25x10 cm (S <sub>2</sub> )	36.67	130.04	7.41	11.92	23.07	34.99	33.91
30x10cm (S₃)	39.26	139.91	8.74	13.22	25.77	38.98	34.11
S. Em±	0.52	1.90	0.24	0.25	0.33	0.63	0.53
CD (P=0.05)	1.52	5.53	0.69	0.72	0.95	1.85	1.56
Fertilizer application							
N:P:K:S 45:35:35:20	35.96	126.71	7.02	11.19	23.02	34.22	32.72
(F <sub>1</sub> )							
N:P:K:S 60:45:45:30	38.20	134.66	8.30	12.62	24.57	37.19	33.95
(F <sub>2</sub> )							
S. Em±	0.43	1.55	0.20	0.20	0.27	0.52	0.44
CD (P=0.05)	1.24	4.52	0.58	0.59	0.78	1.51	1.27
Varieties							
Rcr41 (V <sub>1</sub> )	35.61	128.57	6.26	11.54	23.02	34.56	33.40
Ajmer Coriander ACr <sup>-1</sup>	40.39	135.82	8.11	12.80	24.76	37.56	34.07
(V <sub>2</sub> )							
Pant Haritima (V3)	32.35	123.14	8.62	11.62	23.78	35.40	32.82
S. Em±	0.52	1.90	0.25	0.25	0.33	0.63	0.53
CD (P=0.05)	1.52	5.53	0.72	0.72	0.95	1.85	1.56
	5.95	6.17	13.15	8.82	5.82	7.51	6.80

Table 2. Effect of spacing, fertilizer application and genotypes on yield and yield attributes in coriander

Treatments	Cost of cultivation	Gross returns	Net returns	Benefit : Cost Ratio	
	(Rs. ha <sup>-1</sup> )	(Rs. ha⁻¹)	(Rs. ha <sup>-1</sup> )		
Crop geometry					
20x10 cm (S <sub>1</sub> )	35393	131497	96103	3.72	
25x10 cm (S <sub>2</sub> )	33760	128220	94460	3.80	
30x10cm (S <sub>3</sub> )	32870	125527	92657	3.82	
Fertilizer application					
N:P:K:S 45:35:35:20 (F1)	35657	116923	81267	3.28	
N:P:K:S 60:45:45:30 (F <sub>2</sub> )	37233	126120	88887	3.39	
Genotypes					
Rcr41 (V <sub>1</sub> )	37887	110907	73020	2.93	
Ajmer Coriander ACr <sup>-1</sup> (V <sub>2</sub> )	37887	127160	89273	3.36	
Pant Haritima (V <sub>3</sub> )	35257	117767	82510	3.34	

# Table 3. Economics analysis of different treatments in coriander

Note: Seed selling price @ Rs. 8000 q<sup>-1</sup> prevailing market rate

The data depicted that crop raised at row spacing 30x10 cm (S<sub>3</sub>) recorded significantly higher number of branches (8.86), number of nodes (11.20) at harvest as compared to row spacing of 20x10 cm ( $S_1$ ) and 25x10 cm ( $S_2$ ). Significant improvement in aforesaid parameters due to increase in spacing or in other words reduction in plant population per unit area could be ascribed to availability of more area per plant which implied that individual plant at wider spacing received higher growth inputs (sunlight, water and nutrients) with least competition compared to the plants grown under two closer spacing similar results revealed by the findings of Amin et al.2005 and Narukaet al.2012). Thusgreater inputs at wider spacing resulted in profuse branching which might have helped in larger canopy development and delayed plant to attain reproductive phase. The larger canopy development associated with profuse branching had increased interception, absorption and utilization of solar energy resulting information of higher photosynthates and finally dry matter per plant. A significant improvement in growth with close spacing was in conformity with the findings of Singh and Buttar (2005) and Kumar et al. (2006). It was observed that at row spacing of 30x10 cm (S<sub>3</sub>) various yield attributes i.e. number of umbels per plant, number of umbellate/ plant (139.91) and test weight (8.74 g) highest were recorded in 30x10 cm (S<sub>3</sub>) (Table 2). The maximum value for these estimates were obtained at the row spacing 30x10 cm (S<sub>3</sub>) significantly improved while i.e. days to 50 per cent flowering (60.62) minimum was recorded in 20x10cm (S1), whereas maximum number of umbels per plant (39.26) and number of umbellate/ plant (139.91) and test weight (8.74 g) were recorded in 30x10 cm (S<sub>3</sub>) (Table 2). Significantly highest seed yield (13.22g/ha) and biological yield (38.98 g/ha) were recorded with row spacing of 30x10 cm (S<sub>3</sub>) compared to other spacing (20x10cm ( $S_1$ ) and 25x10cm ( $S_2$ ) cm). The profuse branching seem to have led to greater initiation of flowering and adequate supply of metabolites due to the increase in biomass per plant might have helped in retention of flower thereby greater seed formation and seed growth. Similar findings reported by Moosavi et al. 2012 and Mehta et al.(2013). These results justify that overcrowding of plants at closer spacing significantly reduced growth and yield attributes of the crop but compensated the yield to a certain level.

Effect of fertilizer applications: Significantly higher plant height at 90 DAS and at harvest,

number of nodes and primary branches per plant and minimum days taken to 50% flowering were recorded as a result of higher levels of fertilization F<sub>2</sub>- 60:45:45:30 N:P:K:S ha<sup>-1</sup> (Table 1). It is an established fact that higher levels of fertilization is required for the synthesis of chlorophyll protein. and other organic compounds of physiological significance. Thus, increased level of nitrogen in plant by virtue of its increased availability in the soil medium and there after efficient absorption and translocation in various growth by way of active cell division and elongation resulting in greater plant height number of branches. The nitrogen and application might had resulted in larger canopy development and presumably higher chlorophyll content of leaves and essential oil of seeds, as nitrogen actively participate in its formation with profound influence on photosynthetic efficiency plants, this might had led to higher of accumulation of fresh and dry matter. These findings are in close conformity with those of Ghosh (2009), Nayak et al. (2009), Sharangi et al. (2011) and Diwan et al.(2018). In general, the faster growth of plant with nitrogen evidence from increased biomass per plant at successive stages of crop growth showed better availability of metabolites and nutrients. The present trend of increased in seed yield, straw yield and biological vield of coriander with the application of nitrogen was in close conformity with the findings of Nagheraet al (2000), Tripathi et al. (2001) and Menariaet al.(2007).

# 4. CONCLUSIONS

On the basis of two consecutive years research it has been concluded that plant height was highest at 20x10cm (S1) number of nodes and primary branches/plant significantly increased up to row spacing 30x10 cm (S3) and nitrogen level of 70 kg/ha. Ajmer Coriander ACr<sup>-1</sup> (V2) is far superior not only in growth parameters, but also performed well in terms of yield attributes and ultimately raises the income of the farmers under Malwa Plateau of M.P. It may be recommended for Rabi season cultivation to obtain the maximum seed yield (12.80), net return (89273) and high B:C (3.36).

### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

#### **COMPETING INTERESTS**

Authors have declared that no competing interests exist.

#### REFERENCES

- Amin, A. U., Singh, A., Patel, B. S., & Patel, J. K. (2005). Effect of various row spacing and nitrogen levels on drilled rabi fennel (*Foeniculum vulgare* Mill). Agricultural Science Digest, 25(1), 44-46.
- Department of Agriculture & Farmer Welfare, Gol, Data based report State Level; 3, Area & Production of Horticulture crops for 2023-24 (3rd Advance Estimates), Dtd. 04-06-2024
- Diwan, G., Bisen, B. P., & P. M. (2018). Effect of nitrogen doses and row spacing on growth and seed yield of coriander (*Coriandrum sativum* L.). International Journal of Chemical Studies, 6(4), 2768-2772.
- Diwan, G., Bisen, B. P., Bhooriya, M. S., &Kurrey, V. K. (2018). Effect of nutrient uptake on growth and seed yield of coriander (*Coriandrum sativum* L.) variety Jawahar Dhania-2. *International Refereed Peer Reviewed Indexed Quarterly Journal of Science and Agricultural Engineering*, 8(B), 56-59.
- Ghosh, D. K. (2009). Performance of coriander at different doses of nitrogen fertilizer under coconut-based cropping system. *Indian Coconut Journal*, 51(9), 6-10.
- Hans, R., & Thakral, K. K. (2008). Effect of chemical fertilizers on growth, yield, and quality of fennel (*Foeniculum vulgare Mill*). *Journal of Spices and Aromatic Crops*, 17(2), 134-139.
- Kumar, K., Singh, G. P., Singh, N., & Nehra, B. K. (2006). Effect of row spacing and cycocel on growth and seed yield of coriander (*Coriandrum sativum* L.) cv. Hisar Anand. *Haryana Journal of Horticultural Science*, 35(3&4), 350.
- Malav, N. B., & Yadav, S. N. (1997). Effect of row spacing and level of nitrogen on growth and seed yield of coriander. *Indian Cocoa, Arecanut and Spices Journal*, 21(2), 37-41.
- Mehta, R. S., Meena, S. S., & Vishal, M. K. (2013). Yield and economic feasibility of ajwain (*Trachyspermumammi*) production under varying irrigation interval, nutrient

levels and crop geometry. *Agricultural Science Digest*, 33(1), 56-59.

- Menaria, B. L., &Maliwal, P. L. (2007). Quality of fennel as influenced by plant density, fertilization, and plant growth regulators. *Indian Journal of Plant Physiology*, 12, 57-62.
- Mishra, S. P., Padhiary, A. K., Nandi, A., Patnaik, & Panda, D. (2019). Effect of soluble fertilizer (19:19:19) in coriander (*Coriandrum sativum* L.) (Var. Super Midori). Journal of Pharmacognosy and Phytochemistry, 8(4), 795-797.
- Moosavi, S. G. R., Seghatoleslami, M. J., (2012). &Zareie. Μ. Η. The effect of planting date and plant density on morphological traits and essential oil yield of coriander (Coriandrum sativum L.). International Journal of Agricultural Crop Science, 4, 496-501.
- Naghera, R. P., Sukhadia, N. M., & Ramani, B. B. (2000). Effect of sowing dates and varying levels of nitrogen and phosphorous on coriander (*Coriandrum sativum* L.). *Gujarat Agricultural University Research Journal*, 26(1), 52-54.
- Naruka, I. S., Singh, P. P., Barde, M., & Rathore, S. S. (2012). Effect of spacing and nitrogen levels on growth, yield, and quality of ajwain (*Trachyspermumammi* L. Sprague). *International Journal of Seed Spices*, 2(1), 12-17.
- Nayak, B. R., Saini, S. S., & Sahu, G. S. (2009). Effect of farm yard manure, nitrogen, and plant spacings on yield and attributing characters of coriander (*Coriandrum sativum* L.). *Environmental and Ecology*, 27(3A), 1374-1377.
- Painkara, S. K., Chandrakar, D. K., Paraye, P. M., Verma, A. K., Langangmeilu, G., & Patel, A. (2024). Coriander yield characteristics as influenced by varied date of sowing and planting geometry under Chhattisgarh plain zone. International Journal of Environment and Climate Change, 14(2), 199-204.
- Panse, V. G., &Sukhatme, P. V. (1985). Statistical methods for agricultural workers. ICAR, New Delhi, India.
- Sharangi, A. B., Chatterjee, R., Nanda, M. K., & Kumar, R. (2011). Growth and leaf yield dynamics of cool season coriander as influenced by cutting and foliar nitrogen

Aswani et al.; J. Exp. Agric. Int., vol. 46, no. 12, pp. 581-588, 2024; Article no.JEAI.127684

applicatio	n. <i>Journal</i>	of P	lant Nutrition,
34(12), 17	762-1768.		
Tripathi, A. K.,	Pandya, R.	K., &	Tripathi,M. L.
(2001).	Effect	of	nitrogen,

phosphorous, and potassium on stem gall disease and yield of coriander. *Annals of Plant Protection Sciences*, 9(2), 337-339.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/127684